

VALUING MORTALITY ATTRIBUTABLE TO PRESENT AND FUTURE EXTREME TEMPERATURES IN ARGENTINA

Christian García-Witulski¹, Mariano Rabassa², Mariana Conte Grand³ and Julie Rozenberg⁴

This paper analyzes the effect temperature on human mortality rates in Argentina. The study rests on semiparametric techniques applied to data with a panel structure to estimate the causal effect of temperature extremes on mortality. Monthly mortality rates for municipalities, constructed from the universe of deaths between 2010 and 2019 were regressed on monthly temperatures distributions, precipitation, and municipality-by-month and month fixed effects. Separate regressions were carried out to examine the heterogeneous impacts by age and gender. We also explore alternative specifications as well as differences by death causes. Results show that extreme temperatures increase mortality rates relative to mean temperatures, and the impact of colder-than-average temperatures is slightly greater in magnitude than that of hotter ones. There exists substantial heterogeneity between age groups, with older people facing greater risks while results for gender and proxied income level is inconclusive. All days of extreme cold cause associated death damages equivalent to 0.6% of GDP, while heat damages correspond to 0.1% of 2019 GDP. On the other side, when climate change impacts are valued, damages total 0.7% of GDP in scenario 8.5 since savings due to less mortality occurring in cold day compensate only partially the increase in the number of hot days. If temperature changes are mild (RCP 4.5), mortality would decrease at the national level, but some regions would still be affected.

Keywords: temperature; climate change; mortality; value of a statistical life; Argentina

JEL codes: I18, Q51, Q54

I. Introduction

Climate change represents a major threat to human beings and other forms of life on our planet. The potential impact of climate change on the social and environmental determinants of health outcomes is at the heart of both scientific and policy making efforts. The World Health Organization (WHO) reports that climate change is expected to cause approximately a quarter million additional deaths per year worldwide between 2030 and 2050 due to climate-sensitive diseases, malnutrition, diarrhoea, malaria and heat stress (WHO 2022).

Presently, the global mean temperature is approximately 1.1 degree Celsius above the pre-industrial (1850-1900) era (WMO 2021). There is strong agreement among scientists in the research community regarding the human influence on the observed and predicted changes in temperature and precipitation patterns. Compared to 1850–1900, global surface temperature averaged over 2081–2100 is very likely to be higher by 2.1°C to 3.5°C in the intermediate GHG emissions scenario (SSP2-4.5) and by 3.3°C to 5.7°C under the very high GHG emissions scenario (SSP5-8.5) (Zhongming et al., 2021). Understanding the effects that extreme temperatures could

¹ Facultad de Ciencias Económicas, Universidad Católica Argentina. Corresponding author: christian_garcia@uca.edu.ar

² Facultad de Ciencias Económicas, Universidad Católica Argentina.

³ The World Bank.

⁴ The World Bank.

have on human health is key to promote both effective adaptation policies to rising temperatures from climatic change, and mitigation policies around the planet.

The potential impacts of weather changes on mortality are usually divided in two categories: direct effects (i.e., those produced by temperature increases) and indirect effects arising mainly from changes in vector-borne diseases and food security or increases in weather-related natural disasters. There are in fact other indirect impacts: it is known that weather tends to affect economic growth, energy consumption, agricultural production, labor productivity and pollution levels and all those impacts would likely produce a feedback effect towards worse health (Deschênes & Greenstone, 2011).

Across the literature, some articles limit themselves to study the present sensitivity of mortality at different temperature levels (e.g. (Otrachshenko et al., 2017), for Russia) and others study also temperature mortality aspects due to climate change (e.g., (Deschênes & Greenstone, 2011) or (Barreca, 2012) for the US; (Burgess et al., 2017) for India; or (Yu et al., 2019) for China). The few articles that study climate change impacts on mortality use some of IPCC greenhouse gases emissions' scenarios projections (Representative Concentration Pathways: RCP2.6, RCP4.5, RCP6.0 or RCP8.5⁵) or alternative scenarios. For example, (Yu et al., 2019) consider two extreme pathways, RCP2.6 and RCP8.5, whereas (Burgess et al., 2017) take the Hadley CM3 A1F1 scenario that authors affirm corresponds to a business-as-usual trajectory, and (Deschenes, 2018) follow the CCSME3 A2 scenario.

The causal effect of temperature on mortality rates is usually non-linear. A U-relationship between thermal ambient levels and mortality has been extensively documented in the literature. There are several alternatives to account for nonlinearity: log linear functions (e.g., (Curriero et al., 2002)); polynomial functions (Basu & Samet, 2002); splines, which are piecewise polynomial functions (e.g., (Kaiser et al., 2007); (Doyon et al., 2008); (Barreca, 2012); (Barreca & Shimshack, 2012)); or, bins ((Deschênes & Greenstone, 2011); (Otrachshenko et al., 2017); (Deschenes, 2018); (Yu et al., 2019), among others). Bins arise from discretizing the temperature in parts. If daily temperatures in a month within a year are used, then the marginal effect for a given bin is the additional effect of "one more day" for which the temperature falls into that bin. The main advantage of bins over the other ways of capturing non-linearity is that they offer a very flexible functional form that allow each temperature bin to have a different impact on mortality. Since most of the recent papers in economics use bins, we follow that same approach to ease the comparison.⁶

As highlighted by several authors as (Deschenes, 2014) or (Son et al., 2019), even when we restrict our review to those articles that use bins, it is not easy to compare the results because of several reasons. There are many differences among studies related to: the units of measurement of mortality rates and weather conditions as well as on valuation methodologies.

⁵ For a more detailed description of scenarios characteristics, see (Meinshausen et al., 2011).

⁶ (Deschenes, 2014) (p. 609) states that "The temperature-bins approach is used in most economic papers in the literature, whereas in public health the spline approach is more common." Another difference is that public health literature often uses time-series econometrics instead of individual level regression or panel data. Instead of a time series approach, we use repeated cross-sectional data (i.e., a panel) because we believe this approach might help overcome the problem of relatively short time series usually observed in Argentinian data.

First, in general with respect to deaths, since heat or cold related illnesses are not coded separately under the International Classification of Diseases, studies typically relate all-causes mortality to ambient measures of weather. In addition, some articles use monthly mortality ((A. I. Barreca, 2012), for example), while others use yearly mortality rates ((Deschênes & Greenstone, 2011); (Otrachshenko et al., 2017); (Yu et al., 2019), among others), and only a few use daily mortality (Cohen & Dechezleprêtre, 2019).

Second, in what concerns meteorological conditions, temperature and precipitations are the most usual control variables included in the empirical studies. For temperatures, there is no homogeneous definition of what “cold day” or “hot day” is and so each of the studies divides temperature bins differently.⁷ Since the mortality risk is expected to change with the level of temperature, comparisons lose precision. Another aspect is that most articles use mean temperature (e.g., (A. I. Barreca, 2012); (Deschênes & Greenstone, 2011), among others), and only in few cases, minimum or maximum temperatures are also considered for the regressions (e.g., (Yu et al., 2019) or (Cohen & Dechezleprêtre, 2019)).

Another variable that is sometimes included in mortality-weather regressions is humidity ((A. I. Barreca, 2012)). According to the review performed by (Deschenes, 2014) (p. 609), adding precipitation or humidity “do not lead to meaningful changes in model estimates compared to models that only control for temperatures”. Other authors combine temperature and humidity in a “heat index” to capture “apparent temperature” (Yu et al., 2019), and also find that the change in mortality estimates are “almost negligible”.⁸

Third, once mortality impacts are calculated, a few articles estimate the annual total of years of life lost ((Deschenes & Moretti, 2009); (Burgess et al., 2017); (Cohen & Dechezleprêtre, 2019)). To do that implies leaving aside monetary valuation⁹, and differentiate life years lost depending on each age group life expectancy, but it is not exempted from problems. As (Deschênes & Greenstone, 2011) (p.183) explicitly state, people dying from extreme weather may not be the median citizen and so his/her life expectancy will likely be lower than that of the average citizen. Another thing that may happen is that life expectancies may vary along time and that shift is not considered in this type of calculation.

Moving beyond life-years impacts requires an economic assessment. There are articles that use the present value of foregone earnings of a death among working population, which requires to know deaths by age group, income by age as well as labour force participation (Otrachshenko et al., 2017). This perspective is what is known as the human capital approach, and as such constitutes a lower bound, since productivity lost is only a component of the value of a statistical life, there are also utility losses when a life ends. Hence, present value of lost

⁷ Moreover, some studies use statistical indicators to define a reference temperature to which all the days should be compared, whereas other opt for a definition based on “comfortable temperature”. As discussed in (Otrachshenko et al., 2017) (p. 291), “A typical range of the comfortable temperature limits for a human body is from 68 °F to 74 °F (20–23.3 °C) during the winter and from 73 °F to 78 °F (22.3–25.6 °C) during the summer”.

⁸ A more recent study ((Yuan et al., 2020)) find that humidity has a role to play in determining heat stress, however, even if that was the case of Argentina, we do not have information to include that possibility in our study.

⁹ (Burgess et al., 2017) mentions product and wages losses, but do not combine economic analysis with life years lost.

income neglects the value of leisure time and important subjective aspects and focuses only on pecuniary impacts of death. The full willingness derives from what is known as the Value of a Statistical Life (VSL) estimates.^{10,11} In this case, an option could be to assign the same value of a statistical life -VSL- for all age groups, another would be to calculate the value of a statistical life per year (VSLY), which allow a differentiation of the value of life for people in different age groups since expected remaining years up to life expectancy varies by age group. (Deschênes & Greenstone, 2011) lead these types of estimations since it uses two alternatives: one is to assign a fixed value of a statistical life-year of \$100,000 for the different years lost due to deaths in each age group and the other allows for the value of statistical life-year to be adjusted over time by a real per capita income growth of 2% per year and an elasticity of the value of the statistical life-year to income of 1.6. In any of those cases, the value can come from local data or from a transfer valuation adjusted by relative income and by the elasticity of VSL to income or both.

Although the existing studies use a variety of periods, models, mortality indicators, meteorological variables, the number and size of the weather bins, the reference bins, and ways to value, the great majority of publications find that temperature extremes lead to significant changes in mortality. This assessment can be made more precise if we illustrate that with a few of those articles that use temperature bins with panel data and so are clearly related to our work.

(Deschênes & Greenstone, 2011) is representative of numerous studies for the US on mortality-temperature links. The authors find, for the 1968 to 2002 period, that an additional day with a mean temperature exceeding 90 °F leads to an increase in annual age-adjusted mortality rate of 0.11% per cent with respect to the one within the 50-60 °F interval. Similarly, a day with temperatures below 20 °F is associated with an increase in annual mortality between 0.07 and 0.08 per cent. When they take account for climate change, they estimate that mortality will increase 3 per cent by the end of the century.

(Otrachshenko et al., 2017) analyze the case of Russia in the 1989-2014 period and find that one extra day with temperatures above 25 °C with respect to the 20-25 °C range increase yearly mortality by 0.06%, whereas an additional day with low temperature (below -30 °C) only has no significant impacts.¹² Using the present value of foregone income of working population,

¹⁰ (Schelling, 1968) introduced the idea that valuing risk to life requires capturing individuals' willingness to pay for life-risk reductions (usually referred as the value of a statistical life or VSL). The concept of subjective value of the risk to life is defined as the marginal rate of substitution between risk and consumption: how much consumption the individual is willing to give up reducing his or her risk of dying.

¹¹ Two categories of methods are typically employed to capture VSL: indirect methods (based on revealed preferences) and direct methods (based on stated preferences). The former generally deal with hedonic estimations, relating wages to labor risks (see (Viscusi & Aldy, 2003) for a review). The VSL was more often assessed using hedonic pricing for the labor market ((Viscusi, 1978); (Blomquist, 2004)). The latter method exposes the individuals to hypothetical scenarios of varying risk and cost in which they have to make choices. The idea of capturing the value of life based on directly approaching individuals appeared previously in the work of (Mishan, 1971). The most usual techniques are contingent valuation (CV) and choice experiments (CE). As is the case for hedonic pricing, CV has been extensively applied to health impacts valuation (Cropper et al., 2011).

¹² The authors believe that the result for cold days has to do with a decrease in accidents and diseases' interpersonal transmissions because people, under that type of weather, avoid going outdoors.

the authors estimate that an extra day of heat implies damages equivalent to 0.28% of daily Russian GDP.

(Burgess et al., 2017) study the case of India for the 1957-2000 period and compare it to same results for the US. They find that 1 additional day over 95 °F with respect to the 70-74 °F bin in India increases annual mortality by 0.74% when that increase for the US is of 0.03% for the same years.

(Yu et al., 2019) find for China between 2014-2012, that an extra day of heat (over 90 °F) over the 50-60 °F level increases annual mortality rate by 0.6%, whereas an extra day of cold (less than 10 °F) increases that mortality rate by 0.3%. The authors also show that climate change raises yearly mortality rates in 2.4% and 14.2% in scenarios RCP2.6 and 8.5 respectively. Using the VSL for China, estimated climate change damages account for 0.21% and 1.19% depending on two climate scenario projection.

(Carleton et al., 2020) analyze based on information for 40 countries in the world, that an extra day of heat (above 35°C) increase annual world mortality by 0.05% (4 deaths per million people) whereas an extra cold day (below -5°C) increases that rate by 0.04% (3 deaths per million). The authors also show that if climate change occurs as projected by RCP4.5, mortality could increase by 14 deaths every 100,000 people and 84 deaths/100,000 under RCP8.5. Using the value of a statistical life, those impacts on mortality are equivalent to 0.6 and 3.2 percent respectively of the world GDP. The same authors find that the relationship between temperature and mortality is flattened for high income countries, which can explain part of the differences among studies.

In addition to the overall mortality result, the literature also discusses factors affecting vulnerability to temperature levels ((Hajat & Kosatky, 2010); (Deschenes, 2014); (Benmarhnia et al., 2015); (Son et al., 2019)). One of the clearer ones is age. Elderly are known to face higher risks from cold and heat, which may be due to physiological conditions, housing or health access conditions as well as behavioural characteristics.¹³ Findings suggest higher mortality impact from temperature levels in less qualified employees, among black population and unmarried persons (Son et al., 2019). Other individual-level modifiers as gender or education do not yield conclusive results, whereas lower income individuals tend to be more affected by temperature extremes¹⁴.

The literature analyses other determinants that have to do with community characteristics rather than individual ones. In that sense, there is weak evidence for higher heat related mortality associated with higher population density, poor housing facilities, and higher pollution levels. Additional factors, but also with unsubstantial evidence, are higher risks for heat: in urban areas; in communities with warmer climate; places with less proportion of green areas; etc. Some few studies also report higher vulnerability to cold weather: in regions with

¹³ Despite of that apparent clarity in the results, middle age groups are more affected than the elderly in some cases (this seems to happen for Russia according to (Otrachshenko et al., 2017) (p. 295).

¹⁴ In addition to gender and education, (Cohen & Dechezleprêtre, 2019) also assess how the effect of temperature on mortality changes with income. They find for Mexico that 11% of annual deaths are associated to climate and that temperatures are more likely to kill people in the bottom half of the income distribution. (Barreca et al., 2015) also studies the differences of mortality-temperature impacts across income deciles in the US. They find that cold and hot days impact the most in lowest deciles and the mortality influence of temperature is almost null in the top income deciles. (Carleton et al., 2020) also show the impact differences driven by countries' income.

warmer climate, in places located in lower latitudes and in rural areas (Burgess et al., 2017). In an attempt to take into account those community factors, it is usual to find studies within this literature that test for regional subsamples ((A. I. Barreca, 2012) and (Deschênes & Greenstone, 2011) for different regions in the US; (Yu et al., 2019) for different regions in China; (Carleton et al., 2020) and (Deschenes, 2018) for different countries). From the available literature reviews, it is possible to conclude that, except for the age factor, the rest of the determinants of temperature-mortality sensitivity are location and population sensitive, from where the importance of performing local studies.

This study makes several contributions. Firstly, it adds to the limited existing literature on climatic determinants of mortality in developing countries and in Argentina. A literature review on temperature-related mortality between 1980 and 2017, (Son et al., 2019) shows that most of the studies focus on North America, Europe and Asia, and “many nations, including most in Latin America and Asia, had no studies”. Argentina is an interesting case in its own right because it has a vast territory with diverse weather. The wide range of altitudes, that go from high mountains along the Andes plains on the Atlantic coast, contribute to that diversity.¹⁵ The weather is mild in the Pampas, cold in the western Patagonia, subtropical in the Mesopotamia region, and warm in the Northeast. Along the Andes (located in the West), are the coldest areas, with dry and snowy weather.

Second, it provides estimates of the incidence of climate change on mortality (and not only for the historical link between temperature and death rates). We do so for the whole country (i.e., a sizable portion of the population), breaking the analysis by age and gender groups and using state-of-the-art fixed effects panel data econometrics.

Third, we value mortality damages. In a context of climatic change, where resources must be allocated to promote adaptation to the new climate, comparable countrywide estimates of health impacts and the corresponding damages are indeed needed to develop a rational and sustainable adaptation policy. A few studies have documented the influence of weather on human mortality in Argentina. Of those, the ones that are closer to our work are (Almeira et al., 2016), which reports a U-shaped relationship between temperature and mortality for the cities of Buenos Aires and Rosario, and (de Garín & Bejarán, 2003), that examines the effect of thermal stress during summertime (characterized by the relative strain index) on mortality rates in Buenos Aires city.^{16,17} Although informative, none of the existing

¹⁵ Under the Köppen climate classification, Argentina has 11 different climate types (Rubio, 2018): Humid Subtropical (Cfa, Cwa), moderate oceanic (Cfb), warm semi-arid (BSh), subtropical highland oceanic (Cwb), warm desert (BWh), cold semi-arid (BSk), cold desert (BWk), moderate Mediterranean (Csb), cold oceanic (Cfc), and tundra (ET).

¹⁶ Other studies, are (Chesini, Abrutzky, Herrera, et al., 2019) for the mortality associated to extreme cold in Argentina, (Chesini et al., 2018) and (Chesini, Abrutzky, & Titto, 2019) for heat, but they use very different methods, based on time series instead of on panel data.

¹⁷ Other studies for Argentina focus on the impact of meteorological variables on morbidity. Using data from emergency room visits at a hospital in central Buenos Aires, (Rusticucci et al., 2002) reports the association between weather conditions and several pathologies for both winter and summer times during the 1996-97 season. Yet other studies have turned their focus to specific pathologies. (Alexander, 2013) studies the association between meteorological variables and diverse diseases in the calls to the

work relating weather conditions and health in Argentina perform any economic valuation and this is an important contribution.

The rest of the paper is organized as follows. Section II describes the empirical strategy implemented to value mortality impacts due to temperature variations. Section III describes the data. Section IV shows the estimation results in terms of mortality rate changes, consequent deaths and valuation of those impacts, as well as the robustness checks we perform. The last section V summarizes our findings, the policy consequences they have and discusses the limitations of our results.

II. Empirical strategy

The empirical strategy involves four different steps for the case of Argentina. First, estimate a model that relates daily temperature in a recent period with the corresponding mortality rates. Second, calculate the change in temperature according to climate change projections. Third, estimate deaths attributable to extreme temperatures in the benchmark period as well as the ones resulting from climate change. Fourth, value deaths using the value of a statistical life.

II.1. Benchmark model

Our benchmark model is described by the following equation:

$$MR_{dmta} = \beta_0 + \sum_{j=1}^6 \theta_{aj} \cdot TEM_{dmtj} + \sum_{z=1}^4 \lambda_{az} \cdot PRE_{dmtz} + \mu_m + \omega_{dm} + \varepsilon_{dmta} \quad [1]$$

where MR_{dmta} is the mortality rate per 100,000 inhabitants in municipality or district d (which belong to a given state or province) in month m in year t for a give age-group a . TEM_{dmtj} is the number of days in each district d per month m per year t that fall in each temperature bin j , and PRE_{dmtz} is the number of days in each district d per month m per year t that fall in each precipitation bin j . As in several articles in this literature ((Deschênes & Greenstone, 2011); (Deschenes, 2018)), the temperature range was divided in 10 degrees Fahrenheit (°F) bins¹⁸. The temperature bin taken as the reference is the one of 70-80 °F (≈ 21.1 - 26.6 °C), that corresponds to the mode of the frequency distribution of days in a month that fall into each of the bins. Then, μ_m stands for month time effects that allow controlling for potentially confounding effects common to all municipalities that vary overtime, for example macroeconomic shocks affecting the countrywide economic performance, which might affect health outcomes. Finally, ω_{dm} is an interaction term, which captures temporal variations between municipalities due to differences in air quality, income levels, educational attainment, and population density. Thus, these fixed effects could be interpreted as a baseline estimate of mortality rates in each

public ambulance emergency service of the city of Buenos Aires during the years 1999-2004. (Piccolo et al., 1988) documented the relationship between asthma-related hospitalizations and meteorological variables in the city of Bahia Blanca. More recently, two studies for the city of Córdoba found that a higher daily mean temperature and a wider daily temperature range are important determinants of infectious diseases in both the upper and lower respiratory tracts, particularly among the elderly population and in low-income households enduring housing deprivation ((Amarillo & Carreras, 2012); (Carreras et al., 2015)).

¹⁸ The precipitation range was partitioned in five bins of 4 millimetres (mm) of rain.

municipality ((Barreca & Shimshack, 2012)). Finally, ε_{dmta} is an error term. Note that we use the monthly mortality to avoid the so-called “harvesting effect”: i.e., we try to elude the temporal displacement of deaths that occur that after a period of more deaths than expected because of the heat, follows a period of mortality deficit because it is likely that heat wave especially affected those whose health weak and so would have died in the short-term anyway. If less than a month was considered, we would likely overestimate the impact of temperature on mortality.

Our main estimate for the calculation of damages is in θ_{aj} , which can be read as the change in MR with respect to the MR in the temperature bin of reference, when the number of days in each of the other temperature bins varies by 1:

$$\Delta MR_{dmtaj} = \widehat{\theta}_{aj} \cdot \Delta TEM_{dmtj} , \text{ with } \Delta TEM_{dmtj} = 1 \quad [2]$$

Hence, $\Delta MR_{dmtaj} = \widehat{\theta}_{aj}$

Since the final unit of analysis is municipality (d) by month (m) in a given year (t), there will be a $\widehat{\theta}$ for each age-group (a) for each temperature bin (j): $\widehat{\theta}_{aj}$. More specifically, we ran a model for all age groups and both genders, then, we estimate four regressions, one for each age group. In addition, we run alternative specifications. Finally, we perform some robustness check with the model including temperature and precipitation interactions; using the logarithm of the mortality rate; and using monthly lagged bins.

II.2. Climate change estimation

With the different values for $\widehat{\theta}_{aj}$ obtained from equation [1], we estimate how climate change modifies in each district d the number of days in a given month m of a certain year t that fall within each bin j (ΔTEM_{dmtj}), using climate scenarios for temperature change predictions for the future as compared to the present period. With those two inputs in hand, the change in the mortality rates due to the modification in the number of days that fit within each bin results from:

$$\Delta MR_{dmtaj} = \widehat{\theta}_{aj} \cdot \Delta TEM_{dmtj} \quad [3]$$

More specifically, what is calculated to fulfil the objective of this work is the change in the mortality rates due to the shift in the number of extreme cold (c), heat (h) days, and both extreme types of weather (e), as:

$$\begin{aligned} \Delta MR_{dmtac} &= \widehat{\theta}_{ac} \cdot \Delta TEM_{dmtc} \\ \Delta MR_{dmtah} &= \widehat{\theta}_{ah} \cdot \Delta TEM_{dmth} \\ \Delta MR_{dmtae} &= \widehat{\theta}_{ac} \cdot \Delta TEM_{dmtc} + \widehat{\theta}_{ah} \cdot \Delta TEM_{dmth} \end{aligned} \quad [3']$$

Note that c (for “cold”) corresponds to $j = \text{bin } <40 \text{ }^\circ\text{F}$ ($\approx <4.4 \text{ }^\circ\text{C}$), whereas h (for “hot”) is $j = \text{bin } >90 \text{ }^\circ\text{F}$ ($\approx >32.2 \text{ }^\circ\text{C}$).

II.3. Number of deaths due to extreme temperature and its change

Based on the changes in mortality rate from [2] and [3], we calculate the number of deaths that result from each of three circumstances (changes in cold days, heat days, and both extreme temperature days).

First, we calculate the number of deaths for the whole country resulting from an extra day within each of the bins with respect to the deaths occurring at the reference temperature bin based on equation [2],

$$\Delta Deaths_{aj} = \underbrace{\widehat{\theta}_{aj}}_{\Delta MR_{dmtaj}} \cdot 12 \cdot \frac{Pop_a^{2010}}{100\,000} \cdot n_d \quad [4]$$

Where $\Delta Deaths_{aj}$ is the change in the number of deaths in age-group a for each additional day of temperature within each bin j , $\widehat{\theta}_{aj}$ are the estimated coefficients for average monthly mortality rate from equation [1], 12 are the number of months in a year (this converts monthly mortality to annual mortality), Pop_a^{2010} is the population of group a in 2010 (it is divided by 100 000 because $\widehat{\theta}_{aj}$ is the ΔMR_{dmtaj} in 100 000), and n_d is the number of districts in Argentina (because $\widehat{\theta}_{aj}$ is an average per district that has to be converted to a national estimate, $n_d = 511$ ¹⁹).

Second, to get the change in the number of deaths attributable to each bin temperature per year, we multiply [4] by the average number days in each bin per month during the period used for the estimation of the benchmark model (and then by 12, the number of months in a year):

$$\Delta Deaths_{aj} = \underbrace{\widehat{\theta}_{aj}}_{\Delta MR_{dmtaj}} \cdot TEM_{dmtj} \cdot 12 \cdot \frac{Pop_a^{2010}}{100\,000} \cdot n_d \quad [5]$$

Third, we estimate the number of deaths attributable to the change in the number of days within each bin following equation [3] as:

$$\Delta Deaths_{aj} = \underbrace{\widehat{\theta}_{aj} \cdot \Delta TEM_{dmtj}}_{\Delta MR_{dmtaj}} \cdot 12 \cdot \frac{Pop_a^{2010}}{100\,000} \cdot n_d \quad [6]$$

As it is clear from equations [4]- [6], overall, the total effect of climate change on deaths varies by the population within each age-group in the whole country and depends on the sensitivity of mortality to colder and hotter periods, on the number of cold and hot days in a year, and on how these changes when the distribution of temperature shift as the result of climate change. The economic valuation of such effects is the only step that remains.

II.4. Mortality damages attributable to temperature extremes and their change

Finally, the $\Delta Deaths_{aj}$ calculated in equations [4]- [6] are valued here using the value of a statistical life.

Even when there are non-market methods to assess the VSL as we mentioned above, they involve detailed data and sophisticated techniques, which require resources and

¹⁹ This is an underestimation that may be adjusted in the next version. As of 2020, there is the capital, 135 *partidos* y the Province of Buenos Aires, 380 *departamentos* in the rest of the country. The city Buenos Aires is not included in the results reported here. It will be added in the new version of this document.

knowledge. This may explain why there are relatively few estimates of VSL for low and middle-income countries, as this is widely acknowledged in the literature (see (Robinson, 2017), among others). However, there is an estimation of the VSL in Argentina (Picasso & Grand, 2019). The estimation is not in the context of health but of crime. However, we use it as an alternative because it is obtained with local data. To cover the uncertainty surrounding that applicability of that VSL for this case, we opt for an alternative: transfer values from studies in other countries. Based on them, we adjust VSL from other places (policy site: denoted by p) to Argentina, which is location of the study (s) relative per capita income (I_p/I_s) and the VSL income elasticity (η), as it is usual in the literature (Brouwer, 2000) following:

$$VSL_p = (I_p/I_s)^\eta \cdot VSL_s \quad [7]$$

Then, damages are obtained in each case as the product of $\Delta Deaths_{aj}$ and VSL_p .

III. Data

The first part of our estimation strategy outlined above relies on data for the 2010-2019 period: death counts by administrative subdivision and weather records from meteorological stations Argentinean data. Then, for the second part of our empirical methodology, climate change is captured through the difference between 2010 (the year within the interval that corresponds to the last census done in Argentina) and 2085 distributions of temperature for Argentina as described in the *NASA Earth Exchange Global Daily Downscaled Projections (NEX-GDDP)*.

III.1. Mortality rate

The mortality data correspond to the universe of over 4 million deaths across the country in the period, as compiled by the *Dirección de Estadísticas e Información Pública*, the agency in charge of the *National System of Health Statistics*. The dataset reports each death along with the date of death, the gender and age, and the municipality of residence of the deceased. Thus, death counts were aggregated at the municipal level, the smallest administrative subdivision in the dataset.

Death counts were supplemented with annual population estimates to construct monthly municipal mortality rates per 100,000 inhabitants, by gender and age-groups. Specifically, following (Deschênes & Greenstone, 2011) and (A. Barreca et al., 2015), the population was stratified into four age groups (0-4, 5-44, 45-64, >64-year-olds)²⁰. Annual population numbers by municipality had to be interpolated from the 2010 national population census.

Average mortality rates are (per 100,000) approximately 21 inhabitants for children under 5 years of age, 8 for people between 5 and 44 years of age, 58 for adults between 44 and 65, and 413 for older adults. One should expect to observe the greatest impacts of extreme temperatures in the last group, given that it generally includes the most vulnerable people.


²⁰ Unfortunately, available information did not allow for the construction of separate mortality rates for infants, i.e., younger than one year old, who are known to be far more susceptible to thermal stress than other young children (Keim et al., 2002).

III.2. Meteorological records

Weather records were obtained from the *Servicio Meteorológico Nacional*, the official national weather and climatological agency. Daily maximum and minimum temperatures in degree Celsius (°C) as well as total precipitation in millimetres (mm) are reported for the 71 weather stations that comprise the *National Weather Station Network*. The monitoring network might seem to be sparse given the surface area and the degree of development of a country such as Argentina. However, according to the 2010 national census, the Argentinian population is mostly urban, with 91% living in towns bigger than 2,000 inhabitants, and half of the total population living in just 8 cities. Therefore, the 61 weather stations finally included in the analysis provide coverage to approximately 82% of the country's population.²¹

Daily station data had to be spatially interpolated at the municipal level to construct monthly weather exposure variables. Following a common approach in the literature, square inverse distance weights were calculated from each municipal geographical centroid to each station within a 100 kilometers radius (Hanigan et al., 2006). Thus, the interpolated weather variables are simply the weighted average of the stations' records within that radius (i.e., the closest monitoring station to a particular district is given more weight). Limiting the cut-off distance to 100 km was necessary to balance the number of municipalities included in the study while minimizing the potential measurement error that would otherwise be introduced by including distant stations. Despite districts were not included because of those reasons, they cover 84.9 per cent of Argentina's population. In general, monitoring stations are not located where population is scarce. Similar cut-off levels have been used in empirical estimates for the United States ((Deschênes & Greenstone, 2011); (A. I. Barreca, 2012); (A. I. Barreca & Shimshack, 2012); (Ranson, 2014); (A. Barreca et al., 2015)). Figure 1 show the municipalities included in the study along with the location of the 71 weather stations.

Figure 1. Municipalities included in the study



²¹ This is consider enough given that the paper for China refers to 6% of the population and that for Russia covers 92% of the inhabitants.

Source: Own elaboration.

Notes: The departments in white do not have a meteorological station close at less than 100 km from its center. However, those are places with few inhabitants, and that may be in fact the reason why they do not have a meteorological station close by.

III.3. Climate change

Climate change temperature data comes from the *NASA Earth Exchange Global Daily Downscaled Projections (NEX-GDDP)*. That database has daily minimum and maximum temperature as well as precipitation for the whole planet. The temporal coverage is from 1950 to 2014 historical data and 2015 to 2100 projected Shared Socioeconomic Pathways: RCP-4.5 and RCP-8.5. The RCP 4.5 is a moderate scenario where some effort to curb climate change is made but temperatures rise 2.7°C globally by the end of the century, whereas RCP-8.5 projects that the global economy grows fuelled by exploiting fossil fuels and energy-intensive lifestyles and by 2100 the average global temperature is 4.4°C higher. Note that the temperature by 2050 under the 8.5 scenarios would be similar to that of the 4.5 projection to the end of the century.


The spatial resolution of *NEX-GDDP* is approximately 25 degrees x 25 degrees. We took the daily information for temperature available within the geographical limits of Argentina and average it for both 2010 and 2085.²²

Then, as previously, the temperatures are assigned to each municipality. Figure 2 shows the shift in days within each temperature bin if the projected climate change takes place. What happens in that case is that, as expected, the average temperature increases and the number of cold days decreases while the number of hot days increases.²³ Scenario 8.5, since it projects more climate change, shows a higher number of hot days than the 4.5 one.

²² The geographical references for Argentina are as follows. North: Latitude 21° 46' 52" S, Longitude 66° 13' 17" O; South: Latitude 55° 03' 21" S, Longitude 66° 31' 25" O; East: Latitude 26° 15' 59" S, Longitude 53° 38' 15" O; West: Latitude 50° 01' S, Longitude 73° 34' O.

²³ Note that because NASA data are historical only up to 2014, there is a gap between the base 2010-2019 meteorological information used to estimate the temperature-mortality rate sensitivity and the temperature data from *NEX-GDDP*. This is a usual matter in this type of estimations (Yu et al., 2019). Figure A.1 in Annex shows the gap between our local data and the *NEX-GDDP* model. For example, NASA data does not show days within the > 90 °F while local data does. Other studies correct that gap (Burgess et al., 2017), which we leave for future work.

Figure 2. Change in the distribution of days within a year that fit into each temperature bin in 2010 and 2085 as projected by NASA



Source: Own elaboration.

Note: Average temperature increase 2.2 F in RCP 4.5 and 6.8 F in RCP 8.5. RCP 4.5 at 2085 is approximately RCP 8.5 at 2050. Prediction in 2010 is NASA information, which differs from the national information used to estimate the coefficients of our model. Figure A.1 in the Annex shows that discrepancy.

III.4. Valuation

Finally, for the Value of a Statistical Life, we use an estimation for Argentina from (Picasso & Grand, 2019) that to our knowledge is the only one derived from local data. In addition, for robustness check, we transfer the VSL from estimates in the US (Viscusi & Masterman, 2017) and from a global study by the World Bank in association with the Institute of Health Metric and Evaluation (World Bank & Institute for Health Metrics and Evaluation, 2016).

The local VSL is \$1,5 million corresponding to 2014 (Table 1). The original US estimate is \$9.6 million 2014 dollars, which transferred using the Argentina to US relative GDP per capita yields a value of \$2.4 million.²⁴ Similarly, the World Bank estimate for OECD countries is \$3.8 million, which transferred to Argentina becomes \$1.98 million.

We update the estimates to 2019 dollars, which is the last year of our mortality data, and transfer them to Argentina's case adjusting by the relative GDP ratio among countries. Because of the uncertainty surrounding what is the best VSL, we perform a Montecarlo simulation. To do so, we assume a triangular distribution for VSL, with a minimum value, a maximum value and a mode of 1.2, 1.72 and 1.59 respectively.²⁵

²⁴ For simplicity, in both transfers, we assume that the elasticity of value of a statistical to income is 1. We will also assume uncertainty in last version.

²⁵ Note that the values for 2019 are lower than the original studies because Argentina's per capita GDP has decreased from 2014 to 2019.

Table 1. Value of a statistical life used in our calculations

Reference Country, Year	Picasso and Conte Grand (2019) Argentina, 2014	Viscusi and Masterman (2017) US, 2014	World Bank and IHME (2016) OECD, 2014
	Units	Atlas method current million US\$, 2014	PPP constant 2011 international \$US
Source, previous to any adjustment	1.5	9.6	3.8
Argentina per capita Atlas current million \$US GDP 2014		12,460	
United States per capita Atlas current \$US GDP 2014		55,980	
Argentina GNI per capita, PPP constant 2011 international \$			19,101
GDP per capita Reference for the World Bank results (World Development Report, World Bank)			37,000
Argentina GDP per capita current \$US 2019	12,335		
<i>VSL adjusted</i>	10,006	2.14	1.98
<i>VSL en millones de dólares corrientes de 2019</i>	1.20	1.72	1.59

Source: Own elaboration based on references.


IV. Results

This section is divided in two subsections. The first shows the estimates for the mortality rate temperature impacts for the whole population, by age-groups in the 2010-2019 period. The second subsection uses the estimated relationships to predict the impacts on annual deaths due to temperature extremes, its variation due to climate changes, and their valuation.

IV.1. Mortality rates

The changes in mortality rates attributable to the 2010-2019 temperature profile are summarized in Figure 3. That figure presents the estimated exposure impacts computed by nonparametric regressions between mortality rates and temperature at the municipal level, controlling for precipitation levels and fixed effects by month and municipality-by-month (i.e., equation [1]) for all age-groups, genders, and regions. The results must be interpreted as the relative impact of daily mean temperature over annual mortality due to an additional day in a given month with temperature above or below the observed mode temperature bin during the 2010-2019 period (i.e., the 70-80 °F or 21.1-26.6 °C bin). Any day with a mean daily temperature above the reference temperature positively and significantly affects mortality rates. Days in the extreme cold temperatures (<40 °F bin) also shows a significant excess mortality with respect to the reference temperature.

Figure 3 Impact of temperature on average monthly mortality rates by district



Source: Own elaboration.

Note: The broader 95 confidence intervals are likely due to the fact that there are less observations, so less precision, in the extremes.

The U shape is usual in the literature ((Deschênes & Greenstone, 2011), for example).²⁶ But, the U is not always perfectly symmetric. As Figure 3 suggests, for Argentina, the estimated impacts of colder-than-average temperatures are slightly higher than the computed estimates for hotter than average temperatures. In other countries as Mexico, the higher impacts are also associated to cold and slightly below mean temperature days rather than to heat (Cohen & Dechezleprêtre, 2019).²⁷ More precisely, a cold day (average temperature below 10 °C) cause 3.5 times more excess mortality than a heat day (average temperature above 32 °C).²⁸

The last row of Panel A in Table 2 shows the same result as Figure 3 in a more detailed way. On average, an additional day with a relative cold day, one with a countrywide mean temperature between 30 and 40 °F (<-4.4 °C) increases the population-weighted average monthly mortality rate in 0.633 each 100,000 inhabitants with respect to the mortality for a day with the reference temperature, and that increase is significant. On the other side, an extra relatively hot day (> 90 °F or 32.2 °C) significantly increases that mortality rate by 0.515 in 100,000. If we were to calculate the difference between the curve in Figure 3 and the x axis, excess mortality due to a temperature different from the ideal everyday would be such that the annual mortality rate goes from 586 to 660 per 100,000 in a year.

²⁶ There are also few studies for which the U shape does not appear clearly (this is the case of (Otrachshenko et al., 2017) for Russia, for example).

²⁷ The authors explain that result by the fact that in Mexico just below average mean temperature occur with cold nights and that induces high pathology risks.

²⁸ Note that the change in the U shape for extremely cold days may indicate some adaptive behavior: people may stay home or wear warmer clothes when weather is the coldest and so mortality differentials may be less high than under slightly fewer cold days (see Figure 3).

Table 2. Impact of extreme temperatures on monthly mortality rate per 100,000 inhabitants (2010-2019)

Panel A. Estimates by age group						
°F	bin 40<	bin 40.50	bin 50.60	bin 60.70	bin 80.90	bin > 90
1. 0-4	0.197*** (0.040)	0.161*** (0.029)	0.088*** (0.018)	0.001 (0.021)	0.042 (0.035)	0.112 0.238
2. 5-44	0.008 (0.010)	0.010 (0.006)*	0.001 (0.004)	-0.005 (0.004)	0.040*** (0.007)	0.068* (0.041)
3. 45-64	0.220*** (0.052)	0.367*** (0.040)	0.191*** (0.037)	0.005 (0.024)	0.069* (0.041)	0.304 (0.263)
4. >64	5.566*** (0.484)	5.542*** (0.534)	2.942*** (0.584)	0.841*** (0.124)	0.892*** (0.332)	3.958** (1.877)
5. Population-weighted aggregate estimate	0.633*** (0.069)	0.657*** (0.069)	0.346*** (0.071)	0.084*** (0.022)	0.133*** (0.049)	0.515* (0.288)
Panel B. Alternative specifications						
6. Women	0.645*** (0.108)	0.723*** (0.085)	0.343*** (0.086)	0.068** (0.030)	0.103* (0.061)	0.598 (0.404)
7. Men	0.617*** (0.079)	0.665*** (0.084)	0.384*** (0.079)	0.098*** (0.027)	0.170*** (0.059)	0.492* (0.299)
8. Model with temperature interactions and precipitation	0.687*** (0.103)	0.756*** (0.072)	0.340*** (0.071)	0.107*** (0.023)	0.165*** (0.049)	0.400 (0.271)
9. Model with the logarithm of the mortality rate	0.613*** (0.061)	0.649*** (0.055)	0.427*** (0.029)	0.104*** (0.026)	0.160*** (0.059)	0.542** (0.236)
10. Model with bin lags	0.511*** (0.072)	0.664*** (0.073)	0.365*** (0.091)	0.038 (0.035)	0.002 (0.067)	0.386 (0.423)
11. Model with interaction bin 40< × percentile 10	0.743***† (0.094)	0.653*** (0.070)	0.347*** (0.071)	0.083*** (0.022)	0.133*** (0.049)	0.515* (0.288)
12. Model with interaction bin >90 × percentile 90	0.631*** (0.069)	0.655*** (0.069)	0.343*** (0.071)	0.081*** (0.022)	0.124*** (0.049)	0.807***†† (0.316)

Source: Own calculation

Notes: * p<0.05, ** p<0.01, *** p<0.001. The estimate 5 in panel A corresponds to averages weighted by the population of each age group for the year 2010. The reference BIN is 70-80 °F. All specifications include precipitation bins, municipality and month fixed effects, and province by year fixed effects.

Given that several factors can affect the sensitivity of mortality to temperature levels, regressions were stratified by age-group (Table 2. Panel A), a proxy for income and gender (Table 2. Panel B) in order to examine potential heterogeneity in the temperature-mortality relationship. The elderly is clearly the age-group most affected by temperature since all coefficients are significant in all bins and are much larger than those corresponding to the rest of the age-groups and the population weighted average, as in the literature.²⁹ That impact for elderly is also larger for colder temperatures than for hotter ones. There does not seem to be a pattern regarding gender, which is in line with existing literature. Robustness checks do not show results very different from those for the whole population.

IV.2. Extreme temperatures Deaths and Economic Damages

Once obtained the impact on mortality rates of extreme temperatures in a period and climate change, deaths can be calculated following equations [4]-[6]. The results of those calculations for extreme temperatures are summarized in Table 3.

The last two columns of Panel A in Table 3 show the number of deaths per age group when an additional day of extreme cold and extreme heat occur. They are the result of replacing in equation [4] the coefficients from the first column of Table 1. Overall, the annual countrywide number of deaths due to an extra day of cold is 283 and 229 are the deaths due to an extra day of heat. Those deaths represent 0.072 and 0.070% respectively of Argentina's deaths in 2010 (i.e., 318,477). There are six extreme cold days (less than 4.4. degree Celsius average for the day)

²⁹ A similar result relation between elderly and overall mortality impact of heat is found by (Yu et al., 2019) for China and (Carleton et al., 2020) for a sample of 40 countries, for example.

and two extreme heat days (over 32.2 degree Celsius as the average for a day) in a year. Hence, annual deaths attributable to extreme temperature represent 0.53% and 0.12% of total all cause deaths in Argentina (see 1,767 and 412 deaths in the first two columns of Panel A in Table 2). The latter derive from equation [5]. Note that we are only accounting for excess deaths in the extremes.

Deaths related to changes in extreme temperatures because of climate change are reported on Panel B in Table 3 for the 8.5 and 4.5 scenarios respectively. In the most aggressive scenario in terms of climate change (RCP-8.5), there would be approximately 22 more days of heat and 10 less days of extreme cold per year. Because of the change expected in lower temperature days, mortality decreases in 2,942 deaths, whereas variations in heat days increase mortality by 5,019 cases. Hence, the net effect of temperature changes attributable to climate change accounts for 0.63% of Argentina's yearly average deaths 2010-2019 (i.e., 2,078 per year).

Table 3. Yearly deaths due to extreme cold and heat in Argentina

Panel A. Period 2010-2019

2010-2019 period	Total excess deaths per year		Deaths extra day within bins			
	Due to extreme cold (BIN < 40 °F)	Due to extreme heat (BIN > 90 °F)	extremes	BIN 30 to 40 °F	BIN > 90 °F	
Age range						
0 to 4	46	7	53	7	4	
5 to 44	13	32	45	2	18	
45 to 64	113	45	158	18	25	
> 64	1,595	327	1,922	256	182	
Total deaths	1,767	412	2,179	283	229	
Number of days per year within bins	6	2	8			
% of average 2010-2019 deaths	0.53%	0.12%	0.66%	0.09%	0.07%	

Panel B. With climate change

Age range	Change in total deaths per year			
	Due to extreme cold (BIN < 40 °F)	Due to extreme heat (BIN > 90 °F)	Due to both extremes	
Scenario RCP-8.5				
0 to 4	-76	91	16	
5 to 44	-22	392	370	
45 to 64	-189	550	361	
> 64	-2,655	3,986	1,331	
Total deaths	-2,942	5,019	2,078	
Change in the number of days per year within bins	-10	22		
% of average 2010-2019 deaths	-0.89%	1.51%	0.63%	
Scenario RCP-4.5				
0 to 4	-17	3	-15	
5 to 44	-1	2	1	
45 to 64	-20	7	-12	
> 64	-497	95	-402	
Total deaths	-535	107	-428	
Change in the number of days per year within bins	-2	1		
% of average 2010-2019 deaths	-0.16%	0.03%	-0.13%	

Source: Own elaboration

Note: It has to be remembered that the change in temperature as in NASA does not match exactly the estimated model.

In the less extreme climate change scenario, there would be around 2 days less of extreme cold and 1 day extra (at the maximum) of heat. As a result, mortality would decrease, so there would be gains from climate change. However, there are places where mortality will increase due to climate change since temperature variations are not uniform across the whole country (*Figure A.2 in Annex A*) and inhabitants' resilience may also differ.

Once deaths are accounted for, the economic valuation is straightforward. Table 4 shows the estimated damages. All days of extreme cold cause damages equivalent to 0.5% of GDP, while heat damages correspond to 0.1% of 2019 GDP. On the other side, when climate

change impacts are valued, damages total 0.7% of GDP in scenario 8.5 since saving due to less mortality occurring in cold day compensate only partially the increase in the number of hot days (a similar finding was made for China by (Yu et al., 2019)). In the scenario 4.5 climate change is indeed beneficial. Argentina seems to fit in the middle when compared to those for China (Yu et al., 2019): 0.21% for the RCP2.6 “optimistic” scenario and 1.19% for the “pessimistic” RCP8.5 emissions’ projection path.

Table 4. Damages due to yearly extreme temperature and to expected climate change

	Extreme cold	Extreme heat	Net effect in extremes
Deaths per year			
2010-2019 (local data)	1,767	412	2,179
% total deaths 2010-2019	0.5%	0.1%	0.7%
Number of days per year within bins	6	2	8
Change scenario 8.5 (NASA data)	-2,942	5,019	2,078
Change in the number of days per year	-10	22	
Change scenario 4.5 (NASA data)	-535	107	-428
Change in the number of days per year	-2	1	
Total deaths 2010-2019			331,557
Damages (as % of GDP)			
2010-2019 (local data)	0.6%	0.1%	0.8%
Change scenario 8.5 (NASA data)	-1.0%	1.8%	0.7%
Change scenario 4.5 (NASA data)	-0.19%	0.04%	-0.15%
Argentina GDP million current USD 2019			449,663

Source: Own calculation.


V. Robustness checks

We perform several robustness checks: first, we separate mortality by cause to confirm that cardiovascular and respiratory mortality are the ones that are more closely related to temperature variation; second, we estimate the mortality-temperature relationship by region to assess if people living in regions with low frequencies of cold (hot) days have higher impacts when it is cold (hot) than those which are used to cold (hot) days; third we do try other specifications for the model.

Since the mortality database registers the main cause of death, we compare the mortality-temperature sensitivity for cardiovascular episodes, those associated to respiratory illnesses and malignant neoplasms. As can be seen in Figure 4 cardiovascular related mortality is more responsive to temperatures higher than the reference, and this is particularly the case for high temperatures. Respiratory causes are also sensitive to temperature, but less than cardiovascular-related deaths. This is a usual result in the literature ((Otrachshenko et al., 2017);

(Yu et al., 2019)). Finally, and as expected for the robustness of the results, malignant neoplasm is almost unrelated to the profile of temperatures.³⁰

Figure 4. Excess mortality per cause of death



Source: Own elaboration.

As can be seen in Table 2 Panel B alternative specifications do not change substantially the results.

VI. Conclusions

It is believed that environmental factors are important determinants of human health outcomes. Recently, due to global warming and the recurrence of heatwaves, attention has been placed on how weather affects health outcomes. Of those outcomes it is mortality that has captured the most attention in part because there are some results that show that most of damages of climate change are linked to mortality. According to (Hsiang et al., 2017) approximately 66% of expected damages have to do with mortality in the US: for an increase of average temperature 8 degree Celsius, the cost attributable to mortality is 7% of GDP (when temperature rises around 3 degrees, expected costs are around 1% of GDP).

The existing literature, mainly for developed countries, has pointed to a generally U-shaped type of relationship between temperature and mortality risks. Using flexible regression models over longitudinal monthly weather and mortality data for most municipalities in Argentina spanning seven years, from January 2010 to December 2019, our study has confirmed this relationship from a countrywide perspective since it encompasses 85 per cent of the

³⁰ This same result is found by other authors (see (Deschenes, 2014) p. 614). However, almost not related is not zero related. As shown in (A. I. Barreca, 2012), Panel B Figure 3) cold is somehow positively related to cancer. As discussed in Yu et al (2019), there is evidence that cancer cells may grow faster in cold environments.

country's population. This was achieved using weather data from 71 closest meteorological stations.

Temperature extremes impact Argentina's population already (estimated average yearly damages related to mortality are approximately 0.8% of 2019 GDP), and climate change will cause a net increase in excess deaths due to extreme of 0.7% of GDP under a pessimistic scenario, and a gain of 0.15% of GDP if climate changes is mild. (Carleton et al., 2020) also finds that some countries and regions would benefit from climate change: for example, Europe in both optimistic and pessimistic climate scenarios and the US in the optimistic RCP4.5 scenario.

Our study does have limitations. First, some have to do with impacts that we do not account for. Climate change not only affects mortality through temperature variations but it also induces indirect effects as a higher frequency of natural catastrophes that indirectly imply more deaths. Indirect additional deaths are not accounted for in the present study.

Second, there are several limitations in the data we were able to obtain that are not easy to solve. The available population data did not allow us to construct separate mortality rates for infants, i.e. younger than one year old, which are far more susceptible to thermal stress than other young children, as previously mentioned (Keim et al., 2002). Therefore, the fact that our results show no statistically significant influence of extreme temperatures on children's mortality rates has to be taken with caution. However, even if it were possible to separate a 0-1 years old, a significant result would not necessarily occur. (Yu et al., 2019) did not find it for China, (Otrachshenko et al., 2017) did not see that outcome for Russia, and the same happens in (Deschênes & Greenstone, 2011). The reason may be that, where possible, infants are well protected from extreme temperatures by their parents.

Mortality is obviously an extreme outcome, and thus underestimates the overall effect of extreme temperatures on human health because there are many milder outcomes that may be related to temperature. Unfortunately, there is no detailed countrywide medical database available to estimate both mortality and morbidity impacts.³¹

Third, and somehow related to the previous one, we omit to explicitly account for individual and policy induced averting behaviour. A possible source of underestimation comes from the adoption of precautionary measures, particularly air conditioning, which have a protective effect in extreme temperatures (A. Barreca et al., 2015). Adaptation is possible through changes in outdoor activities, geographical mobility, heating and air conditioning adoption or more energy consumption due to a more intensive use of existing cooling and heating appliances. Adaptation may vary by income since it changes access to better healthcare and to heating and air conditioning appliances (Cohen & Dechezleprêtre, 2022). Some articles make an effort to consider adaptation using data for defensive investments as residential energy consumption or air-conditioning acquisition ((Deschênes & Greenstone, 2011); (A. Barreca et al., 2015); (Carleton et al., 2020)), but we do not have that type of data at the county level (to match with our database). Hence, considering explicitly adaptation is out of the scope of our paper. Hence, in that sense, our results can be read as an upper bound of mortality but not of costs since adaptation involves expenditures.

³¹ Morbidity data was requested from the authorities but not submitted apparently because of problems in data quality.

Fourth, we made some assumptions that could be changed. An implicit assumption is that population distribution across municipalities would remain constant under climate change. Mortality changes as a result of climate change depends on population changes in level and in terms of its composition as an aging population implies higher costs since older people are more sensitive to temperature extremes. When considering population aging, the gains from climate change under the RCP4.5 decrease but do not revert their sign.

Fifth, given the sparse distribution of the station network, interpolation of weather parameters from distant stations might have introduced measurement error. To limit the chance of this type of error happening, we follow the convention of excluding weather stations further than 100 kilometres away from each municipal geographical centre. Satellites might provide better weather information, at least in terms of spatial coverage (Kloog et al., 2014), but we leave this to future research. As pointed out before, there is also a gap in the merge between local and international temperature data.

Having pointed out the potential shortcomings of this research, we believe this study provides valuable information on weather determinants of mortality in Argentina. The estimation approach employed in this paper helps to disentangle the causal effect of temperatures on mortality risk while controlling for possible confounders, thus reducing the chance that the observed temperature-mortality link could have been driven by omitted factors associated both with weather and mortality.

Finally, the results indicate that temperature is important in terms of mortality, but is also relevant in terms of the implied monetary damages. Having that information is key for both adaptation and mitigation policies' decisions based on benefit-cost analysis.

Acknowledgements

The authors acknowledge the *Dirección de Estadísticas e Información en Salud* of the Argentina's Ministry of Health for making available daily mortality from 2010 to 2019 to undertake this research. We thank Maria Eugenia Bonilla-Chacin and the Human Development team at the World Bank Buenos Aires office for their comments.

Funding

This work was supported by the World Bank through the Argentina Climate and Development Project (P176901).

References

- Alexander, P. (2013). Association of monthly frequencies of diverse diseases in the calls to the public emergency service of the city of Buenos Aires during 1999–2004 with meteorological variables and seasons. *International Journal of Biometeorology*, 57(1), 83–90.
- Almeira, G. J., Rusticucci, M. M., & Suaya, M. (2016). Relación entre mortalidad y temperaturas extremas en Buenos Aires y Rosario. *Meteorológica*, 41.

- Amarillo, A. C., & Carreras, H. A. (2012). The effect of airborne particles and weather conditions on pediatric respiratory infections in Cordoba, Argentina. *Environmental Pollution*, *170*, 217–221.
- Barreca, A., Clay, K., Deschênes, O., Greenstone, M., & Shapiro, J. S. (2015). Convergence in adaptation to climate change: Evidence from high temperatures and mortality, 1900-2004. *American Economic Review*, *105*(5), 247–251.
- Barreca, A. I. (2012). Climate change, humidity, and mortality in the United States. *Journal of Environmental Economics and Management*, *63*(1), 19–34.
- Barreca, A. I., & Shimshack, J. P. (2012). Absolute humidity, temperature, and influenza mortality: 30 years of county-level evidence from the United States. *American Journal of Epidemiology*, *176*(suppl_7), S114–S122.
- Basu, R., & Samet, J. M. (2002). An exposure assessment study of ambient heat exposure in an elderly population in Baltimore, Maryland. *Environmental Health Perspectives*, *110*(12), 1219–1224.
- Benmarhnia, T., Deguen, S., Kaufman, J. S., & Smargiassi, A. (2015). Vulnerability to heat-related mortality. *Epidemiology*, *26*(6), 781–793.
- Blomquist, G. C. (2004). Self-protection and averting behavior, values of statistical lives, and benefit cost analysis of environmental policy. *Review of Economics of the Household*, *2*(1), 89–110.
- Brouwer, R. (2000). Environmental value transfer: State of the art and future prospects. *Ecological Economics*, *32*(1), 137–152.
- Burgess, R., Deschenes, O., Donaldson, D., & Greenstone, M. (2017). Weather, climate change and death in India. *University of Chicago*.
- Carleton, T. A., Jina, A., Delgado, M. T., Greenstone, M., Houser, T., Hsiang, S. M., Hultgren, A., Kopp, R. E., McCusker, K. E., & Nath, I. B. (2020). *Valuing the global mortality consequences of climate change accounting for adaptation costs and benefits*. National Bureau of Economic Research.

- Carreras, H., Zanobetti, A., & Koutrakis, P. (2015). Effect of daily temperature range on respiratory health in Argentina and its modification by impaired socio-economic conditions and PM10 exposures. *Environmental Pollution*, 206, 175–182.
- Chesini, F., Abrutzky, R., Herrera, N., Skansi, M. de los M., Fontán, S., Gonzalez Morinigo, E., Savoy, F., & De Titto, E. (2019). Mortalidad asociada a eventos extremos de frío en Argentina, 2005-2015. *Revista Argentina de Salud Pública*, 10(41), 28–36.
- Chesini, F., Abrutzky, R., & Titto, E. de. (2019). Mortalidad por olas de calor en la Ciudad de Buenos Aires, Argentina (2005-2015). *Cadernos de Saúde Pública*, 35, e00165218.
- Chesini, F., Herrera, N., Skansi, M. de los M., González Morinigo, É. C., Fontán, S., Savoy, F., & Titto, E. de. (2018). *Análisis de la mortalidad durante las olas de calor del verano 2013-2014 en la República Argentina*.
- Cohen, F., & Dechezleprêtre, A. (2019). Mortality, temperature, and public health provision: Evidence from Mexico. *American Economic Journal: Economic Policy*.
- Cohen, F., & Dechezleprêtre, A. (2022). Mortality, temperature, and public health provision: Evidence from Mexico. *American Economic Journal: Economic Policy*, 14(2), 161–192.
- Cropper, M., Hammitt, J. K., & Robinson, L. A. (2011). Valuing mortality risk reductions: Progress and challenges. *Annu. Rev. Resour. Econ.*, 3(1), 313–336.
- Curriero, F. C., Heiner, K. S., Samet, J. M., Zeger, S. L., Strug, L., & Patz, J. A. (2002). Temperature and mortality in 11 cities of the eastern United States. *American Journal of Epidemiology*, 155(1), 80–87.
- de Garín, A., & Bejarán, R. (2003). Mortality rate and relative strain index in Buenos Aires city. *International Journal of Biometeorology*, 48(1), 31–36.
- Deschenes, O. (2014). Temperature, human health, and adaptation: A review of the empirical literature. *Energy Economics*, 46, 606–619.
- Deschenes, O. (2018). Temperature Variability and Mortality: Evidence from 16 Asian Countries. *Asian Development Review*, 35(2), 1–30.

- Deschênes, O., & Greenstone, M. (2011). Climate change, mortality, and adaptation: Evidence from annual fluctuations in weather in the US. *American Economic Journal: Applied Economics*, 3(4), 152–185.
- Deschenes, O., & Moretti, E. (2009). Extreme weather events, mortality, and migration. *The Review of Economics and Statistics*, 91(4), 659–681.
- Doyon, B., Bélanger, D., & Gosselin, P. (2008). The potential impact of climate change on annual and seasonal mortality for three cities in Québec, Canada. *International Journal of Health Geographics*, 7(1), 1–12.
- Hajat, S., & Kosatky, T. (2010). Heat-related mortality: A review and exploration of heterogeneity. *Journal of Epidemiology & Community Health*, 64(9), 753–760.
- Hanigan, I., Hall, G., & Dear, K. B. (2006). A comparison of methods for calculating population exposure estimates of daily weather for health research. *International Journal of Health Geographics*, 5(1), 1–16.
- Hsiang, S., Kopp, R., Jina, A., Rising, J., Delgado, M., Mohan, S., Rasmussen, D., Muir-Wood, R., Wilson, P., & Oppenheimer, M. (2017). Estimating economic damage from climate change in the United States. *Science*, 356(6345), 1362–1369.
- Kaiser, R., Le Tertre, A., Schwartz, J., Gotway, C. A., Daley, W. R., & Rubin, C. H. (2007). The effect of the 1995 heat wave in Chicago on all-cause and cause-specific mortality. *American Journal of Public Health*, 97(Supplement_1), S158–S162.
- Keim, S. M., Guisto, J. A., & Sullivan, J. B. (2002). Environmental thermal stress. *Annals of Agricultural and Environmental Medicine*, 9(1).
- Kloog, I., Nordio, F., Coull, B. A., & Schwartz, J. (2014). Predicting spatiotemporal mean air temperature using MODIS satellite surface temperature measurements across the Northeastern USA. *Remote Sensing of Environment*, 150, 132–139.
- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C., & Riahi, K. (2011). The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Climatic Change*, 109(1), 213–241.

- Mishan, E. J. (1971). Evaluation of life and limb: A theoretical approach. *Journal of Political Economy*, 79(4), 687–705.
- Otrachshenko, V., Popova, O., & Solomin, P. (2017). Health consequences of the Russian weather. *Ecological Economics*, 132, 290–306.
- Picasso, E., & Grand, M. C. (2019). The Value of the Risk to Life in the Context of Crime. *Journal of Benefit-Cost Analysis*, 10(2), 178–205.
- Piccolo, M., Perillo, G., Ramon, C., & DiDio, V. (1988). Outbreaks of asthma attacks and meteorologic parameters in Bahia Blanca, Argentina. *Annals of Allergy*, 60(2), 107–110.
- Ranson, M. (2014). Crime, weather, and climate change. *Journal of Environmental Economics and Management*, 67(3), 274–302.
- Robinson, L. A. (2017). Estimating the Values of Mortality Risk Reductions in Low-and Middle-Income Countries¹. *Journal of Benefit-Cost Analysis*, 8(2), 205–214.
- Rusticucci, M., Bettolli, L. M., & de Los Angeles Harris, M. (2002). Association between weather conditions and the number of patients at the emergency room in an Argentine hospital. *International Journal of Biometeorology*, 46(1), 42–51.
- Schelling, T. C. (1968). The life you save may be your own. *Problems in Public Expenditure*, 127–162.
- Son, J.-Y., Liu, J. C., & Bell, M. L. (2019). Temperature-related mortality: A systematic review and investigation of effect modifiers. *Environmental Research Letters*, 14(7), 073004.
- Viscusi, W. K. (1978). Labor market valuations of life and limb: Empirical evidence and policy implications. *Public Policy*, 26(3), 359–386.
- Viscusi, W. K., & Aldy, J. E. (2003). The value of a statistical life: A critical review of market estimates throughout the world. *Journal of Risk and Uncertainty*, 27(1), 5–76.
- Viscusi, W. K., & Masterman, C. J. (2017). Income elasticities and global values of a statistical life. *Journal of Benefit-Cost Analysis*, 8(2), 226–250.
- World Bank & Institute for Health Metrics and Evaluation. (2016). *The cost of air pollution: Strengthening the economic case for action*.

Yu, X., Lei, X., & Wang, M. (2019). Temperature effects on mortality and household adaptation: Evidence from China. *Journal of Environmental Economics and Management*, 96, 195–212.

Yuan, J., Stein, M. L., & Kopp, R. E. (2020). The evolving distribution of relative humidity conditional upon daily maximum temperature in a warming climate. *Journal of Geophysical Research: Atmospheres*, 125(19), e2019JD032100.

Zhongming, Z., Linong, L., Xiaona, Y., Wangqiang, Z., & Wei, L. (2021). *AR6 Climate Change 2021: The Physical Science Basis*.


[WHO] World Health Organization. 2022. Climate change and health. Available from: <https://www.who.int/news-room/fact-sheets/detail/climate-change-and-health>. Accessed on 17 February 2022.

[WMO] World Meteorological Organization. 2021. The State of the global climate 2021. World Meteorological Organization. Available from: <https://public.wmo.int/en/media/press-release/state-of-climate-2021-extreme-events-and-major-impacts>. Accessed on 17 February 2022.


Annex

Figure A.1. Gap between Argentina’s meteorological system temperature and NEX-GDDP data

Panel A. Error in Scenario 8.5




Panel B. Error in Scenario 8.5



Note: Present (observed) is the information from the Argentina’ meteorological system whereas Present (prediction) is the NEX-GDDP data.

Figure A.2. Changes in temperature per district: RCP 8.5



Source: Own elaboration.

Note: We include here only the changes in temperature and not in deaths and damages because the number of observations does not allow calculating coefficients per bin per district, and we were to use the national estimates to each district, estimates would not do a good job when compared with actual mortality data.